

---

## FIRST OBSERVATION OF *CHELON LABROSUS* (RISSO, 1827) IN UM- HUFAYAN LAGOON, LIBYA

---

Mohammad El-Sayed El-Mor <sup>1\*</sup>, Amani Fitori Ali<sup>2</sup>, and Esam M. Buzaid <sup>3</sup>

<sup>1</sup> Dept. of Marine Resources, Fac. of Sci., Univ. of Suez Canal, Egypt.

<sup>2</sup> Dept. of Marine Resources, Fac. of Natural Resources, Univ. of Tubruk, Tubruk, Libya.

<sup>3</sup> Dept. of Marine Sciences, Fac. of Sci., Omar Al-mukhtar Univ., Albayda, Libya.

Corresponding author: melmor88@yahoo.com,

---

### ABSTRACT

A presence of *Chelon labrosus* (Risso, 1827) was recorded for the first time in Um-Hufayan Lagoon, Libya; between November 2017 to January 2018. Samples ranged between 11.1 and 25.9 cm in length, and ranged between 25.594 and 166.647 g in weight. The fish's morphometric characters and meristic counts were recorded. The relationships between total length and other morphometric measurements and meristic counts, in addition to length – weight relationship and condition factors were studied.

Results showed that the descriptive traits were matched with the previously recorded *Chelon labrosus* description, and there are some insignificant differences in meristic counts. Also, results revealed that the growth is non-isometric between length and weight of this species. The relationships between length and morphometric traits were statistically non-significant, indicating that they do not change as fish grow.

**Keywords:** *Chelon labrosus*, Um Hufayan lagoon, Eastern Libya

### 1. INTRODUCTION

It cannot be hidden that some rare species have recently been noticed on the eastern coast of Libya, and its shores and estuaries, especially if they have an economic importance. [1] stated that grey mullets are recognized easily from other fishes by the two widely separated small dorsal fins (4 slender spines in anterior fin and a soft-rayed posterior fin). This family is cumbersome to identify, because of the high morphological similarities shared between species, especially in juveniles [2].

Mugilids family includes 64 moderate-sized inshore pelagic species in all tropical and temperate seas. Eight species in the Mediterranean, all in the eastern basin. Some mugilid species often occur unexpectedly in new regions [3 & 4], including the thick-lipped grey mullet *Chelon labrosus* (Risso 1827) in the North Atlantic till the West African coast, and to southern Norway according to [5], and in the Mediterranean Sea and the western Black Sea as well [6]. *Chelon labrosus* is characterized distinctively by its depth as pupil diameter; four spines in first dorsal fin, meanwhile the second dorsal fin has a single spine and 8 rays; thick upper lip and 9 rays in anal fin [7, 8, 9 and 10].

These inshore pelagic fishes swim along the shore, in schools. Spawning season in the winter, and feeding on great variety of food types [11]. Juveniles feed on zooplankton; adults feed on algae, vegetal detritus and sediment [2]. According to [12], females grow faster than males, in the Mediterranean thick-lipped grey mullet get mature sexually at 35.5 cm, and maximum reported length was 60.0 cm [5]. This euryhaline species enters estuaries, lakes and protected shores [10 & 11], caught by trammel net and purse seine in Libyan eastern coast. [13] mentioned that this species annually appears rarely in Susah coast in November.

In general, morphometric and meristic methods remain the simplest and most direct way among methods and techniques to identify fish species [14-24]. [25] mentioned that descriptive traits are non-measurable non countable body morphological characters, and they often vary clinically along geographic gradients [26]. Meanwhile the external measurements of an organism are called morphometric, while meristic is serial counts of body elements.

[27] stated that length-weight relationships are important to: (i) incorporate fisheries and growth models, (ii) to predict weight from length measurements made in the yield assessment, and

(iii) to indicates the robustness and the general well-being of fishes [28]. It is influenced by sex, gonad maturity, health of the fish, seasonal effect, degree of stomach fullness and preservation techniques [29]. The condition factor in fish contributes to understand its life cycle and to adequate management, hence, to maintain the equilibrium in the ecosystem [30 & 31].

This study aims to characterize and identify a new-appearing Mugilidae specimen from Um Huffayan lagoon, using the taxonomic reference collection, and to compare the metric and meristic characters of this species, to that determined in this work.

## 2. MATERIAL AND METHODS

### 2.1. Study site (Um-Hufayan: 32° 33' 13.5" N, 23° 05' 57.2" E):

This location in Bumba bay, about 80 km east of Darna towards to Tobruk, covers an area of about two Km<sup>2</sup> and depth 0.5 - 3m (**Fig. 1**), through it sea water enter the lagoon on tide times. Underground springs at the inner side of the Lagoon discharge water at the inner side of the Lagoon and the salinity may be as low as about 11‰. This ecosystem of this lagoon gave it ability as resting site for migratory birds and nesting site for sea-turtles [32]. Also, this area is important as nursery and feeding ground for many commercial fishes; such as those of the family Cichlids, Sparids, Mugilids and Serranids [32 - 36].

### 2.2. Fish identification and morphometrics:

The fish specimens (**Fig. 1**) were caught in Um-Hufayan Lagoon between November 2017 and January 2018, and taken to the Marine Biology Laboratory of Omar Al-Mukhtar University for identification according to specific literatures [5, 9, 10, 11, 37, 38 & 39].

**2.2.1. Descriptive traits:** [body shape, body color, strips color, shape of mouth and fins].

**2.2.2. Meristic traits:** [Dorsal fin (DF), Pectoral fin (PF), Pelvic fin (PvF), Anal fin (AF), Caudal fin (CF) and Lateral line scales (LL)] according to [13, 39 & 40].

**2.2.3. Total length (TL) – morphometric traits relationships:** TL with [Standard Length

(SL), Forked Length (FL), Body Depth (BD), Head Length (HL), Cheek Length (CL), Eye Diameter (ED), Pre-orbital Length (PoL), Post-orbital Length (PoOL), Pre-Anal length (PAL), Pre-dorsal length(PDL), Pre-pelvic Length (PPL), Mouth Width (MW) and Total Weight (TW)(gm.)], following [24].

### 2.3. Length-weight relationship (LWR): [41- 46]

$$W = aL^b$$

[W= total weight, L= total length,  
a = Constant value,  
b = Variable value (2 – 4)].

### Condition factors of Fulton: [45 & 47]

$$K_F = 100 W_t L^{-3}$$

[W<sub>t</sub> = total weight (g),  
L = total length (cm)].

### Condition factors of Clark: [45 & 48]

$$K_c = 100 W_g L^{-3}$$

[W<sub>g</sub> = gutted weight (g),  
L = total length (cm)].

**2.4. Statistical studies:** Statistical analyses were done by MS Excel 2010, using power equation


## 3. RESULTS AND DISCUSSION

### 3.1. Descriptive traits:

According to the descriptive traits, the spindle-shaped and elongated body, with strongly flattened head, with two short dorsal fins; and thick upper lip and its depth as pupil diameter. Most body is shiny and silvery, upper-side is dark-gray with an olive tinge; sides are light-gray, with elongated and dark stripes. Silvery abdomen, with a whitish shade and dark gray fins, Most of these distinctive characters are matching with the recorded descriptions [5, 10 & 11].

### 3.2. Meristic traits:

Taxonomical formulae [D1: IV; D2: I + 7-10; A: III + 9; P: 17; Pv: I +5; LL: 41-46] is typical meristic characters of *C. Labrosus* specimen (**Table, 1**) according to [5 & 10]. As expected, differences between individual meristic



**Fig. (1):** Um-Hufayn lagoon, Bomba bay, eastern Libya, and *Chelon labrosus* (Risso 1827) individuals.

parameters and fish length were very weak and insignificant for fish indicating that meristic traits are conservative and do not change as fish grows; hence meristic formulae based on these traits can be used to identify and characterize different fish species, although some errors are more likely to increase at high sample size [24 & 49].

### 3.3. Total length-Morphometric traits relationships:

Samples of *C. labrosus* were ranged from  $12.0 \pm 3.5$  to  $24.8 \pm 1.1$  cm in length and from  $34.547 \pm 11.212$  to  $172.531 \pm 86.710$  g in weight. Results showed that morphometric parameters were also related to fish length by power equations (Table, 1 and Fig. 2 A & B). These results showed insignificant relationships between total length and most morphometric traits of specimen of *C. labrosus* from Um-Hufayn, these results are mainly close to the results of [24] on *L. aurata* at the same study area. As well as, all the relationships were negative, indicating that traits increase insignificantly with increase in length, except Pre-dorsal Length (PDL) (2.8221).

### 3.4. Length-weight relationship (LWR)

The length weight relationship of *C. labrosus* was expressed by:  $W = 30.61 L^{1.0068}$  (Table, 1 and Fig. 2A). The constant  $b$  of *C. labrosus* was not close to 3, the value that

indicated that the growth is unisometric. These results are mainly close to results of [24] about *Liza aurata* at the same study area.

### 3.5. Condition factors of Fulton & Clark

Fulton and Clark condition factors were close to 1. In this study, relatively high magnitude of Fulton condition factor ( $K_F$ ) was dropped from  $1.99 \pm 0.59$  (in size group of 12.0 cm) to  $0.87 \pm 0.08$  (in size group of 24.8 cm). Meanwhile, Clark condition factor ( $K_C$ ) ranged from  $1.46 \pm 0.49$  (in the smallest specimens) to  $0.63 \pm 0.05$  (in the largest ones) as shown in Figure (2 B). These average values of condition factors [ $K_F = 1.22$  and  $K_C = 0.89$  (Table 1)] were higher than that recorded for *L. aurata* in the same location [24]. Correlations of  $K_F$  and  $K_C$  with total length of *C. labrosus* were statistically insignificant, indicating that they do not change as fish grow. [50] stated that growth is strongly influenced by both biotic and abiotic environmental conditions, and as index of status of the aquatic ecosystem in fish spam.

### 3.6. Total length frequency distribution

As shown in Fig (2 B), mode of stock was recognized in size group of 21.6 cm (41.9 %), and this zenith for juveniles has larger size than that was recorded for *Liza carinata* samples in Al-Hamamah coasts [55]. Generally, these juveniles are in critical sizes, where sexual maturity begins [56].

**Table (1) . Morphometric and meristic traits (related to fish total length by power equations) of *C. labrosus* from Um-Huffayn, East of Libya, between November 2018 and January 2019.**

Morphometric Parameter	Size groups (cm)					a	b	R <sup>2</sup>
	11.1 - 14.0	14.1 - 17.0	17.1 - 20.0	20.1 - 23.0	23.1 - 26.0			
Size Groups (average) (cm.)	12.0	15.8	18.5	21.6	24.8	-	-	-
Count	3	5	6	13	5	-	-	-
Total Length (TL) (cm.)	12 ± 0.9	15.8 ± 0.8	18.5 ± 1.3	21.6 ± 1.3	24.8 ± 1.2	-	-	-
Standard Length (SL) (cm.)	10.2 ± 1.0	13.7 ± 0.8	16.8 ± 0.8	19.4 ± 1.1	22.6 ± 0.6	9.9998	0.487	0.994
Forked Length (FL) (cm.)	11.0 ± 0.9	14.6 ± 0.8	17.5 ± 1.1	20.5 ± 1.0	23.8 ± 0.7	10.755	0.4709	0.991
Pre-orbital Length (POL)(cm.)	1.1 ± 0.3	0.9 ± 0.2	1.1 ± 0.4	1.2 ± 0.2	1.2 ± 0.4	1.0028	0.091	0.2423
Post-orbital Length(PoOL)(cm.)	1.7 ± 0.5	1.5 ± 0.8	1.9 ± 0.4	2.1 ± 0.7	2.1 ± 0.6	1.5653	0.1716	0.5665
Body depth (BD) (cm.)	3.6 ± 1.3	3.4 ± 0.4	3.9 ± 1.4	4.3 ± 1.2	4.3 ± 1.6	3.4114	0.1352	0.6682
Pre-anal Length (PAL) (cm.)	2.9 ± 0.7	3.8 ± 2.0	3.8 ± 1.3	4.1 ± 1.3	4.4 ± 1.9	2.9925	0.2395	0.9281
Pre-pelvic Length(PPvL) (cm.)	7.2 ± 0.9	6.4 ± 0.6	6.6 ± 0.8	7.6 ± 1.0	7.3 ± 0.6	6.7968	0.0316	0.0775
Pre-dorsal Length (PDL) (cm.)	7.2 ± 1.5	8.8 ± 2.0	9.1 ± 1.4	9.9 ± 1.4	11.3 ± 1.2	0.0175	2.8221	0.9881
Head Length (HL) (cm.)	4.4 ± 0.8	4.1 ± 0.6	3.9 ± 0.4	4.5 ± 0.8	5.0 ± 0.7	4.1261	0.0586	0.1537
Cheek Length (CL) (cm.)	1.4 ± 0.2	1.5 ± 0.5	1.5 ± 0.3	1.6 ± 0.5	1.7 ± 0.7	1.385	0.1085	0.8779
Caudal Peduncle Length (CPL)	0.1 ± 0.6	0.09 ± 0.4	1.9 ± 0.2	1.5 ± 0.8	0.8 ± 0.6	0.119	0.097	0.374
Caudal Peduncle Width (CPD)	2.1 ± 0.2	2.4 ± 0.9	2.6 ± 0.7	2.3 ± 0.4	2.2 ± 0.5	0.141	0.828	0.401
Mouth Gape (MG) (cm)	1.3 ± 0.1	0.4 ± 0.6	1.2 ± 0.7	1.1 ± 0.3	1.9 ± 0.4	0.709	0.318	0.228
Mouth Width (MW) (cm.)	1.7 ± 0.5	1.5 ± 0.8	2.4 ± 0.4	1.9 ± 0.7	2.1 ± 0.6	0.851	0.595	0.223
Eye Diameter (ED) (cm.)	0.7 ± 0.2	0.7 ± 0.1	0.8 ± 0.2	0.9 ± 0.2	1.0 ± 0.1	0.6562	0.2223	0.8127
Dorsal Fin Length (DFL) (cm)	6.2 ± 1.4	6.8 ± 2.0	6.4 ± 1.2	7.6 ± 1.2	6.3 ± 1.2	0.540	0.593	0.429
Anal fin Length (AFL) (cm)	2.6 ± 0.5	3.4 ± 2.1	3.6 ± 1.3	3.4 ± 1.4	4.6 ± 1.2	0.240	1.185	0.604
Pectoral Fin Length (PFL) (cm)	0.8 ± 0.3	0.5 ± 0.1	0.3 ± 0.9	0.4 ± 0.7	0.9 ± 0.1	1.71	0.172	0.430
Snout Length (SL) (cm)	1.3 ± 0.5	1.1 ± 0.2	1.2 ± 0.6	1.9 ± 0.2	1.5 ± 0.1	1.316	0.726	0.646
Pelvic Fin Length (PvFL) (cm)	1.0 ± 0.5	0.6 ± 0.2	1.5 ± 0.6	1.3 ± 0.1	1.5 ± 0.5	0.734	0.665	0.184
Total Weight (TW) (g)	34.5 ± 11.2	53.1 ± 20.8	79.92 ± 13.5	131.5 ± 56.6	172.1 ± 86.7	30.609	1.0068	0.9572
Condition factor – Fulton (K <sub>F</sub> )	1.99 ± 0.59	1.27 ± 0.36	1.00 ± 0.14	0.97 ± 0.19	0.87 ± 0.08	0.227	1.004	0.64
Condition factor – Clark (K <sub>C</sub> )	1.46 ± 0.49	0.92 ± 0.33	0.73 ± 0.14	0.71 ± 0.37	0.63 ± 0.05	0.295	0.864	0.182
Dorsal fin 1 (D1)	IV	IV	IV	IV	IV	0.290	0.971	0.758
Dorsal fin 2 (D2)	I + (8 ± 1)	I + (8 ± 1)	I + (8 ± 2)	I + (9 ± 1)	I + (8 ± 2)	0.089	0.277	0.812
Anal fin (A)	III + (9)	III + (9)	III + (9)	III + (9)	III + (9)	0.317	0.815	0.941
Pectoral fin (P)	14 ± 2	16 ± 1	17 ± 0	17 ± 0	17 ± 0	0.852	0.452	0.317
Pelvic fin (Pv)	I + (5)	I + (5)	I + (5)	I + (5)	I + (5)	0.904	0.354	0.548
Lateral line (LL)	43 ± 2	42 ± 3	44 ± 2	44 ± 3	45 ± 3	0.671	0.931	0.538
Gill rakers	58 ± 9	62 ± 9	65 ± 10	61 ± 10	59 ± 8	1.352	0.825	0.643





Fig (2-A): Meristic and Morphometric traits and total length relationships (related to fish total length by power equations), with condition factors and length frequency distribution of *C. labrosus* from Um-Huffayn, East of Libya, between November 2018 and January 2019.



**Fig (2-B): Meristic and Morphometric traits and total length relationships (related to fish total length by power equations), with condition factors and length frequency distribution of *C. labrosus* from Um-Huffayn, East of Libya, between November 2018 and January 2019.**

#### 4. CONCLUSION

The recent record of *C. labrosus* in this area, is probably evidence that species is highly adaptable to new environmental conditions, meaning that occurrence of this species is significantly related to its euroyhaline nature and a wide temperature range (4–37°C), and great variety of food menu [10, 12 & 51]. As well as, this expansion of rare species into wider ranges, might be related to climate change or human activity [36], Mugilids prompt to enter bays and lower reaches of the Mediterranean brackish waters [10, 52, 53 & 54]. According to [32], Um Huffayn is almost exposed to immigration of juvenile eels and optimum ecosystem for tilapias and mullets. It is essential to perform a detailed characteristic of the new recorded specimen, which may constitute a threat to the fish-diversity in this lagoon.

#### 5. REFERENCES

- [1] Menezes, N.A. ; de Oliveira, C. and Nirchio, M. (2010). Anoldtax on omicdilemma: the identity of the western south Atlanticle branche mullet (Teleostei :Perciformes: Mugilidae). *Zootaxa*, 2519: 59—68.
- [2] Kottelat, M. & Freyhof, J. (2007). *Hand book of European Fresh water Fishes*. Kottelat, Cornol and Freyhof, Berlin, 646 pp.
- [3] Mohra, H. (1988). Zur Biologie und fischereilichen Bedeutung der Meeräsche an der deutschen Nord- und Ostseeküste. On the biology of the Grey Mulletts and their importance for fisheries on the German coast, Protokolle zur Fischereitechnik, Inst. Fangtech., Hamburg.
- [4] Nehring, S. (2002). Biological invasions into German waters: an evaluation of the importance of different human-mediated vectors for non in diagnose macro zoo-benthic species. In: Leppäkoski, E., Golasch, S., Olenin, S. (Eds.), *Invasive Aquatic Species of Europe: Distributions, Impacts and Management*. KluwerAcad. Publ., Dordrecht, 373—383.
- [5] Ben-Tuvia, A. (1986). Mugilidae, [in:] *Fishes of the North-eastern Atlantic and Mediterranean*. Vol. 3, P.J.P. Whitehead, J.C. Bauchot, J.C. Hureau & E Tortonese (eds.), UNESCO, Paris, 1197–1204.
- [6] Keith, P. & Allardi, J. (2002). *Atlas des poissons d’eau douce de France*, Patrimoines Naturels, 47, 387 pp.
- [7] Whitehead, P. J. P.; Bauchot, J. C.; Hureau, J. C.; Nielsen, J. and Tortonese, E. (1984). *Fishes of the North eastern Atlantic and Mediterranean*. Published by the United Nations Educational, Scientific and Cultural Organization. United Kingdom. UNESCO.
- [8] Calypso, (2000). *Mediterranean Fishes. A Modern Taxonomic Checklist. A Listing of all species recorded on the Calypso. Ichthyological Database for the Mediterranean Area*, including all Lessepsian migrants. Calypso Publications. London. UK.
- [9] Nelson, J.S. (2006). *Fishes of the World*, 4<sup>th</sup>ed. John Wiley & Sons, Inc., 601pp.
- [10] Golani, D.; Ozturk, B. and Basusta, N. (2006). *The Fishes of the Eastern Mediterranean*. Turkish Marine Research Foundation, Istanbul, Turkey. 259 pp.
- [11] Ben-Abdalla, A. R.; Al-Gmati, H.; Kasim, A. A.; Al-Turkie, A. A. and Ben-Moussa, M. N. (2009). *Guide to bony fishes in Libyan waters*, Marine biology Research Center (MBRC) - Tajoura’a.
- [12] Harrison, I. J. (2003). *Chelonlabrosus* (Risso, 1826), [in: *Freshwater Fishes of Europe*. Vol. 8, Part I: Mugilidae, Atherinidae, Blenniidae, Odontobutidae, Gobiidae 1, P. J. Miller (ed.), AULA-Verlag, Wiebelsheim, 17–24.
- [13] Rasheed, E. R. (2012). Some ecological and biological studies on some species of family Mugilide inhabiting Susa coast, El-Gabal Al-Akadar, Libya, MSc. Thesis, Fac. of Sci. Zool Dept. Omar El-Mukhtar Univ. pp. 122.
- [14] Ihssen, P.; Booke, H.; Casselman, J.; Mc Glade, J.; Payne, N. and Utter, (1981). Stock identification: Materials and methods. *Can. J. Fish. Aquat. Sci.*, 38: 1838-1855.
- [15] Creech, S. (1992). A multivariate morphometric investigation of *Atherina boyeri* Risso, 1810 and *A. presbyter* Cuvier 1829. (Teleostei: Atherinidae): morphometric evidence in support of the two species. *J. Fish Biol* 41:341-353.
- [16] Mamuris, Z.; Apostolidis, A.P.; Panagiotaki, P.; Theodorou, A.J. and Triantaphyllidis, C. (1998). Morphological variation between red mullet populations in Greece. *J. Fish Biol.*, 52:107-117.
- [17] Bronte, C.R.; Fleischer, G.W.; Maistrenko, S.G. and Pronin, N.M. (1999). Stock structure of Lake Baikalomulas determined by whole body morphology-. *J. Fish Biol.*, 54:787-798
- [18] Swain, D. & Foote, C. (1999). Stocks and chameleons the use of phenotypic variation in stock identification. *Fish Res*, 43; 113-128.
- [19] Zafar, M.; Nazir, A.; Akhtar, N.; Naqvi, S. and Zia-ur-Rehman, M. (2002). Studies on meristic counts and morphometric measurements of Mahseer (Torputitora) from spawning ground

- of Himalayan Foot-hill River korang Islamabad, Pakistan. Pakistan J. Biol. Sci., 5 (6): 733-735.
- [20] Barriga-Sosa, D.; Jimenez-Badillo, M.; Ibanez, A. and Arredondo-Figueroa, J. (2004). Variability of Tilapias (*Oreochromis* Spp.) introduce in Mexico: morphometric, meristic and genetic characters. J. Appl. Ichthyol., 20: 7-14.
- [21] Doherty, D. & McCarty, T. (2004). Morphometric and meristic characteristics analysis of two western Irish populations of Arctic char, *Salvelinus alpinus* (L.). Biology and Environment. proceedings of the Royal Irish Academy, 104:75-85.
- [22] Naesje, T.; Vuorinen, J. and Sandlund, O. (2004). Genetic and morphometric differentiation among sympatric spawning stocks of white fish (*Coregonus uslavaretus* L.) in Lake Femund, Norway. J. Limnol., 63: 233-243.
- [23] Sokal, M. & Oktener, A. (2009). Biometry of the fishes *Barbus plebejus* and *Barbus capito* from Coruh Basin, Turkey. Rev. Biol. Trop, 57:159-165.
- [24] Abdrabba, I. A. A. (2009). Comparative studies on otolith of *Tilapia zillii* and *Liza aurata* from Umm Hufayn lagoon (Gulf of Bumba), Eastern Libya, Omar AL-Mukhtar University, Dept. Of Zool. Faculty of Science,– Marine Biology Branch, Non-published thesis, 95 pp.
- [25] Talwar, P. & Jhingran, A. (1992). Inland fishes of India. Rec. Ind. J., 3: 19-24.
- [26] Lindsey, C. (1988). Factors controlling meristic variation. In Fish physiology, Vol. XI-B (Hoar W S, Randall D J, eds), pp.197-274 .San Diego, CA: Academic Press.
- [27] Nehemia, A.; Justin, D.; Maganira and Cyrus Rumisha (2012). Length Weight relationship and condition factor of tilapia species grown in marine and freshwater ponds. Agriculture and biology J. of north America, ISSN print: 2151-7517.
- [28] Beyer, J. (1987). Length-weight relationships part1: Computing the mean weight of the fish of a given length class Fish byte 5:11-13.
- [29] Tesch. F. W. (1971). Age and growth. In: methods for assessment of fish production in fresh waters. W.E. Ricker (Ed). Blackwell Scientific publications, Oxford: 98-130.
- [30] Imam, T.; Bala, U.; Balarabe, M. and Oyeyi, T. (2010). Length-weight relationship and condition factor of four fish species from Wasai Reservoir in Kano, Nigeria. African Studies.6:125-130.
- [31] Ndimele, P.; Kumolu-Johnson, C.; Aladetohun, N. and Ayorinde, O. (2010). Length-weight relationship, condition factor and dietary composition of Sarotherod on *melano theron* Ruppell, 1852 (Pisces: Cichlidae) in Ologe Lagoon, Lagos, Nigeria. Agric. Biol. J. North Am., Vol. 1: 584-590.
- [32] Abdalhamid, A. H. A.; Ali, R. A. S.; El- Mor, M. E. ; Ali, S. M. and Elawad, A. N. (2018). Some Ecological and Biological Studies on the European eel *Anguilla anguilla* (Linnaeus, 1758) in Umm Hufayan brackish lagoon, eastern Libya Mediterranean Sea. Bulletin de l'Institut Scientifique, Rabat, Section Sciences de la Vie, n° 40: 23-30.
- [33] Lemoalle, J. & Saad, A. (1987). Lagune cotiere de Libya et deltas de Egypt. Pages 63-74 In: African wet land sand shallow water bodies, Burgis M.J. and J. J Symons (eds), Orstom.
- [34] MBRC (2005). Atlas of the Mediterranean Sea, Marine Biology Research Center (MBRC), Tajura, Tripoli, 135 pp.
- [35] Reynolds, J. E.; Haddoud, D. A. and Vallet, F. (1995). Prospects for aquaculture development in Libya, Libfish Field documents No 9. Tripoli / Rome, FAO.
- [36] Badalamenti, F.; Ben Amer, I.; Dupuy De La Grandrive, R.; Foulquie, M., Milazzo, M.; Sghaier, Y.R.; Gomei, M. and Limam, A. (2011). Scientific field survey report for the development of Marine Protected Areas in Libya, 32 pages.
- [37] Whitehead, P. J. P.; Bauchot, J. C.; Hureau, J. C.; Nielsen, J. and Tortonese, E. (1984). Fishes of the North eastern Atlantic and Mediterranean. Published by the United Nations Educational, Scientific and Cultural Organization. United Kingdom. UNESCO.
- [38] Brylińska, M. ( 2000). Rybysłodkowodne Polski, PWN, Warszawa, 521 pp.
- [39] Czerniejewski, P.; Keszka, S. and Rybczyk, A. (2008). *Chelon labrosus* (Risso, 1827) – the first record from Lake Dąbie (Poland), Ocean ologia, 50: 281 – 284.
- [40] Fischer, W.; Bauchot, M.-L. and Schneider, M. (1987). Fiches FAO d'identification des espèces pour les besoins de la pêche.(Révision 1). Méditerranée et mer Noire, Zone de pêche 37. Volume II. Vertébrés, Publication préparée parla FAO, résultat d'un accord entre la FAO et la Commission des Communautés Européennes (Projet GCP/INT/422/EEC), financée conjointement par ces deux organisations. Rome, FAO, Vo1.2: 761- 1530 (In French).
- [41] Hile, R. (1936). Age and growth of the ciscoe, *Leveichthyes artedi* (lesueur), in the lakes of the northern high lands, Wisconsin . Bull.Mar.fish ,U.S., 48(19): 211-317.
- [42] Beckman, W. C. (1948). The weight –length relationship factors of conversion between standard and total lengths and coefficient of condition for seven Michigan fishes. *Trans. Amer. Fish. Soc.*75: 237-256.
- [43] Le Cren, E. (1951). The length-weight relationship and seasonal cycling weight and

- condition in the perch (*Perca fluviatilis*). J. Animal Ecology. 20: 201-219.
- [44] Ricker, W. E. (1975). Computation and interpretation of biological statistics of fish populations. Bulletin of fisheries research board of Canada 191.382 pp.623-625.
- [45] Bagenal, T. & Tesch, F. (1978). Methods of Assessment of Fish Production in Fresh Waters .IBPH and book No3, 3rded. Oxford Black well Scientific Publication, London.101-136.
- [46] Khan, M. A.; Khan, S. and Miyan, K. (2012). Studies on Length-weight and Length-length Relationships of Four Freshwater Fishes Collected from River Ganga. Journal of Fisheries and Aquatic Science, 7: 481-484.
- [47] Fulton, F. (1902). Rate of growth of sea fishes. Sci. Invest. fish. Div. Scot. Rep. 20.
- [48] Clark, F. N. (1928). The weight length relationship of the California Sardine (*Sardina coerulea*) at San-Pedro, *Fish. Bull.* 12: 34-39.
- [49] Mohammed, W. E.( 2018). Morphological Study to identify tilapia species (Cichlidae) in Ain Um-Hufayan Lagoon, Eastern Libya Coast, MSc. Thesis, Fac. of Sci. Zool Dept. Omar El-Mukhtar Univ. pp. 122.
- [50] Anene, A. (2005). Condition factor of four Cichlid species of a man-made Lake in Imo State, South-eastern Nigeria. *Turkish J. Fisheries and Aquatic Sciences*, 5: 43-47.
- [51] Chervinski J. (1977). Adaptability of *Chelon labrosus* (Risso) and *Liza saliens* (Risso) (Pisces, Mugilidae) to fresh water, *Aquaculture*, 11 (1): 75–79.
- [52] Vieira, J.P. ( 1991). Juvenile mullets (Pisces: Mugilidae) in the Estuary of Lagoados Patos, RS, Brazil. *Copeia*, Vol. 2: 409—418.
- [53] Carvalho, C.D.; Corneta, C.M. and Uieda, V.S. (2007). Schooling behavior of *Mugilicurema* (Perciformes: Mugilidae) in an estuary in south-eastern Brazil. *Neotrop. Ichthyol.* 5 (1): 81—83.
- [54] Kottelat, M. & Freyhof, J. (2007). Hand book of European Fresh water Fishes. Kottelat, Cornol and Freyhof Berlin, 646 pp.
- [55] Ekwella, S.A. (2008). Ecological and biological studies on commercial catch in Al-Hammamah coast, eastern Libya. MSc. Thesis, Omar Almkhtar University, 180 pp.
- [56] Akyol, O. (2003). Age, growth and reproduction period of grass goby (*Zosterisessor ophiocephalus* Pallas 1811) in the Bay of Izmir (Aegean Sea). *Archive of Fishery and Marine Research* 50(2):220-224

## الرصد الأول لأسماك البوري *Chelon labrosus* (Risso, 1827) في بحيرة أم حفين، ليبيا

محمد السيد المر<sup>1</sup> أماني فيتوري علي<sup>2</sup> ،

عصام محمود بوزيد<sup>3</sup>

- 1 قسم علوم البحار، كلية العلوم جامعة قناة السويس مصر  
2 قسم الموارد البحرية، كلية الموارد الطبيعية وعلوم البيئة، جامعة طبرق، ليبيا  
3 قسم علوم البحار، كلية العلوم ، جامعة عمر المختار، البيضاء، ليبيا

### الملخص العربي

تم تسجيل ظهور سمك البوري الكيلوني من النوع *Chelon labrosus* لأول مرة في منطقة أم حفين، ليبيا في الفترة من نوفمبر 2017 إلى يناير 2018، حيث تراوحت أطوال العينات بين 11.1 - 25.9 سم في الطول وأوزانها بين 25.594 - 166.647 جم. كما تم تسجيل القياسات التشكلية والصفات العددية لديها. وقد تمت دراسة العلاقة بين الطول الكلي والقياسات التشكلية والصفات العددية الأخرى، بالإضافة إلى تعيين العلاقة بين الطول والوزن للأسماك واستنباط معامل الحالة لها.

وقد أوضحت النتائج تطابق وصف العينات مع الصفات المسجلة سابقاً لهذا النوع، وأن الاختلافات طفيفة وغير معنوية أحصائياً في الصفات العددية. كما أوضحت النتائج أن نمو هذه الأسماك غير متمائل ما بين الطول والوزن؛ وأن العلاقة بين طول الجسم والصفات التشكلية غير معنوية احصائياً مما يدل على أن هذه الصفات لا تتغير مع زيادة الطول.